

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

$I, J, P$  need confirmation.

In the quark model,  $\Xi_b^0$  and  $\Xi_b^-$  are an isodoublet ( $usb, dsb$ ) state; the lowest  $\Xi_b^0$  and  $\Xi_b^-$  ought to have  $J^P = 1/2^+$ . None of  $I, J$ , or  $P$  have actually been measured.

## $\Xi_b$ MASSES

### $\Xi_b^-$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>5797.0 ± 0.6 OUR AVERAGE</b>	Error includes scale factor of 1.7. See the ideogram below.		
5796.70 ± 0.39 ± 0.23	AAIJ	19AB LHCb	$pp$ at 7, 8 and 13 TeV
5797.72 ± 0.46 ± 0.31	1 AAIJ	14BJ LHCb	$pp$ at 7, 8 TeV
5793.4 ± 1.8 ± 0.7	2 AALTONEN	14B CDF	$p\bar{p}$ at 1.96 TeV
5774 ± 11 ± 15	3 ABAZOV	07K D0	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5795.8 ± 0.9 ± 0.4	4 AAIJ	13AV LHCb	Repl. by AAIJ 19AB
5796.7 ± 5.1 ± 1.4	5 AALTONEN	11X CDF	Repl. by AALTONEN 14B
5790.9 ± 2.6 ± 0.8	6 AALTONEN	09AP CDF	Repl. by AALTONEN 14B
5792.9 ± 2.5 ± 1.7	7 AALTONEN	07A CDF	Repl. by AALTONEN 09AP

<sup>1</sup> Reconstructed in  $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ ,  $\Xi_c^0 \rightarrow p K^- K^- \pi^+$  decays. Reference  $\Lambda_b^0$  mass 5619.30 ± 0.34 MeV from AAIJ 14AA.

<sup>2</sup> Uses  $\Xi_b^- \rightarrow J/\psi \Xi^-$  and  $\Xi_c^0 \pi^-$  decays.


<sup>3</sup> Observed in  $\Xi_b^- \rightarrow J/\psi \Xi^-$  decays with  $15.2 \pm 4.4^{+1.9}_{-0.4}$  candidates, a significance of 5.5 sigma.

<sup>4</sup> Measured in  $\Xi_b^- \rightarrow J/\psi \Xi^-$  decays.

<sup>5</sup> Measured in  $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$  with  $25.8^{+5.5}_{-5.2}$  candidates.

<sup>6</sup> Measured in  $\Xi_b^- \rightarrow J/\psi \Xi^-$  decays with  $66^{+14}_{-9}$  candidates.

<sup>7</sup> Observed in  $\Xi_b^- \rightarrow J/\psi \Xi^-$  decays with  $17.5 \pm 4.3$  candidates, a significance of 7.7 sigma.



### $\Xi_b^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>5791.9 ±0.5 OUR AVERAGE</b>			
5794.3 ±2.4 ±0.7	AAIJ	14H LHCb	$pp$ at 7 TeV
5791.80±0.39±0.31	<sup>1</sup> AAIJ	14Z LHCb	$pp$ at 7, 8 TeV
5788.7 ±4.3 ±1.4	<sup>2</sup> AALTONEN	14B CDF	$p\bar{p}$ at 1.96 TeV
••• We do not use the following data for averages, fits, limits, etc. •••			
5787.8 ±5.0 ±1.3	<sup>3</sup> AALTONEN	11X CDF	Repl. by AALTONEN 14B

<sup>1</sup> Uses  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  and  $\Xi_c^+ \rightarrow p K^- \pi^+$  decays. The measurement comes from the mass difference of  $\Xi_b^0$  and  $\Lambda_b^0$ .

<sup>2</sup> Uses  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  decays.

<sup>3</sup> Measured in  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  with  $25.3^{+5.6}_{-5.4}$  candidates.

### $m_{\Xi_b^0} - m_{\Lambda_b^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>177.5 ±0.5 OUR AVERAGE</b>			Error includes scale factor of 1.6.
177.73±0.33±0.14	<sup>1</sup> AAIJ	17BE LHCb	$pp$ at 7, 8 TeV
176.2 ±0.9 ±0.1	<sup>2</sup> AAIJ	13AV LHCb	$pp$ at 7 TeV
••• We do not use the following data for averages, fits, limits, etc. •••			
177.08±0.47±0.16	<sup>3</sup> AAIJ	17BE LHCb	$pp$ at 7, 8 TeV
178.36±0.46±0.16	<sup>4,5</sup> AAIJ	14BJ LHCb	$pp$ at 7, 8 TeV

<sup>1</sup> Combination of the original statistically independent measurements of AAIJ 14BE and AAIJ 17BJ taking into account correlation between systematic uncertainties.

<sup>2</sup> Reconstructed in  $\Xi_b^0 \rightarrow J/\psi \Xi^-$  decays.

<sup>3</sup> Reconstructed in  $\Xi_b^- \rightarrow J/\psi \Lambda K^-$  decays. Reference decays  $\Lambda_b^0 \rightarrow J/\psi \Lambda$  were used.

<sup>4</sup> Reconstructed in  $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ ,  $\Xi_c^0 \rightarrow p K^- K^- \pi^+$  decays. Reference  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ .

<sup>5</sup> Combined with AAIJ 17BE.

### $m_{\Xi_b^0} - m_{\Lambda_b^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>172.5 ± 0.4 OUR AVERAGE</b>			
174.8 ± 2.4 ± 0.5	AAIJ	14H	LHCB $pp$ at 7 TeV
172.44 ± 0.39 ± 0.17	<sup>1</sup> AAIJ	14Z	LHCB $pp$ at 7, 8 TeV

<sup>1</sup> Uses  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  and  $\Xi_c^+ \rightarrow p K^- \pi^+$  decays.

### $m_{\Xi_b^-} - m_{\Xi_b^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>5.9 ± 0.6 OUR AVERAGE</b>			
5.92 ± 0.60 ± 0.23	<sup>1</sup> AAIJ	14BJ	LHCB $pp$ at 7, 8 TeV
3.1 ± 5.6 ± 1.3	<sup>2</sup> AALTONEN	11X	CDF $p\bar{p}$ at 1.96 TeV

<sup>1</sup> Reconstructed in  $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ ,  $\Xi_c^0 \rightarrow p K^- K^- \pi^+$  decays. Uses  $m(\Xi_b^0) - m(\Lambda_b^0) = 172.44 \pm 0.39 \pm 0.17$  MeV from AAIJ 14Z.

<sup>2</sup> Derived from measurements in  $\Xi_b^- \rightarrow \Xi_c^+ \pi^-$  and  $\Xi_b^- \rightarrow J/\psi \Xi^-$  from AALTONEN 09AP taking correlated systematic uncertainties into account.

## $\Xi_b$ MEAN LIFE

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <https://hflav.web.cern.ch/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

### $\Xi_b^-$ MEAN LIFE

VALUE ( $10^{-12}$ s)	DOCUMENT ID	TECN	COMMENT
<b>1.572 ± 0.040 OUR EVALUATION</b>			
<b>1.57 ± 0.04 OUR AVERAGE</b>			Error includes scale factor of 1.1.
1.599 ± 0.041 ± 0.022	<sup>1</sup> AAIJ	14BJ	LHCB $pp$ at 7, 8 TeV
1.55 $^{+0.10}_{-0.09}$ ± 0.03	<sup>2</sup> AAIJ	14T	LHCB $pp$ at 7, 8 TeV
1.36 ± 0.15 ± 0.02	AALTONEN	14B	CDF $p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.56 $^{+0.27}_{-0.25}$ ± 0.02	<sup>3</sup> AALTONEN	09AP	CDF Repl. by AALTONEN 14B

<sup>1</sup> Reconstructed in  $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ ,  $\Xi_c^0 \rightarrow p K^- K^- \pi^+$  decays. Reference  $\Lambda_b^0$  lifetime  $1.479 \pm 0.009 \pm 0.010$  ps from AAIJ 14U.

<sup>2</sup> Measured in  $\Xi_b^- \rightarrow J/\psi \Xi^-$  decays.

<sup>3</sup> Measured in  $\Xi_b^- \rightarrow J/\psi \Xi^-$  decays with  $66^{+14}_{-9}$  candidates.

**$\Xi_b^0$  MEAN LIFE**

VALUE ( $10^{-12}$ s)	DOCUMENT ID	TECN	COMMENT
-----------------------	-------------	------	---------

**1.480 ± 0.030 OUR EVALUATION**

<b>1.477 ± 0.026 ± 0.019</b>	<sup>1</sup> AAIJ	14Z	LHCB $pp$ at 7, 8 TeV
------------------------------	-------------------	-----	-----------------------

<sup>1</sup> Uses  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  and  $\Xi_c^+ \rightarrow p K^- \pi^+$  decays. The measurement comes from the value of relative lifetime of  $\Xi_b^0$  to  $\Lambda_b^0$ .

 **$\Xi_b^-$  MEAN LIFE**

VALUE ( $10^{-12}$ s)	DOCUMENT ID	TECN	COMMENT
-----------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.48 <sup>+0.40</sup> <sub>-0.31</sub> ± 0.12	<sup>1</sup> ABDALLAH	05c	DLPH $e^+ e^- \rightarrow Z^0$
1.35 <sup>+0.37+0.15</sup> <sub>-0.28-0.17</sub>	<sup>2</sup> BUSKULIC	96T	ALEP $e^+ e^- \rightarrow Z$
1.5 <sup>+0.7</sup> <sub>-0.4</sub> ± 0.3	<sup>3</sup> ABREU	95v	DLPH Repl. by ABDALLAH 05c

<sup>1</sup> Used the decay length of  $\Xi^-$  accompanied by a lepton of the same sign.

<sup>2</sup> Excess  $\Xi^- \ell^-$ , impact parameters.

<sup>3</sup> Excess  $\Xi^- \ell^-$ , decay lengths.

 **$\tau_{mix}$  (1/2 $\pi$ ) times the oscillation period**

VALUE (s)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

$> 13 \times 10^{-12}$	<sup>1</sup> AAIJ	17BH	LHCB $pp$ at 7, 8 TeV
------------------------	-------------------	------	-----------------------

<sup>1</sup> Uses  $\Xi_b^{*-}$  and  $\Xi_b^{\prime-}$  decays to  $\Xi_b^0 \pi^-$ , where  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$ ,  $\Xi_c^+ \rightarrow p K^- \pi^+$ .

**MEAN LIFE RATIOS** **$\tau_{\Xi_b^-} / \tau_{\Lambda_b^0}$  mean life ratio**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>1.089 ± 0.026 ± 0.011</b>	<sup>1</sup> AAIJ	14BJ	LHCB $pp$ at 7, 8 TeV
------------------------------	-------------------	------	-----------------------

<sup>1</sup> Reconstructed in  $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ ,  $\Xi_c^0 \rightarrow p K^- K^- \pi^+$  decays. Reference  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ .

 **$\tau_{\Xi_b^-} / \tau_{\Xi_b^0}$  mean life ratio**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>1.083 ± 0.032 ± 0.016</b>	<sup>1</sup> AAIJ	14BJ	LHCB $pp$ at 7, 8 TeV
------------------------------	-------------------	------	-----------------------

<sup>1</sup> Reconstructed in  $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ ,  $\Xi_c^0 \rightarrow p K^- K^- \pi^+$  decays. Uses  $\Xi_b^0$  measurements from AAIJ 14Z.

 **$\Xi_b^-$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\Xi^- \ell^- \bar{\nu}_\ell X \times B(\bar{b} \rightarrow \Xi_b^-)$	$(3.9 \pm 1.2) \times 10^{-4}$	S=1.4
$\Gamma_2$ $J/\psi \Xi^- \times B(b \rightarrow \Xi_b^-)$	$(1.02^{+0.26}_{-0.21}) \times 10^{-5}$	
$\Gamma_3$ $J/\psi \Lambda K^- \times B(b \rightarrow \Xi_b^-)$	$(2.5 \pm 0.4) \times 10^{-6}$	
$\Gamma_4$ $p D^0 K^- \times B(\bar{b} \rightarrow \Xi_b^-)$	$(1.7 \pm 0.6) \times 10^{-6}$	

$\Gamma_5$	$\rho \bar{K}^0 \pi^- \times B(\bar{b} \rightarrow \Xi_b^-)/B(\bar{b} \rightarrow B^0)$	$< 1.6$	$\times 10^{-6}$	CL=90%
$\Gamma_6$	$\rho K^0 K^- \times B(\bar{b} \rightarrow \Xi_b^-)/B(\bar{b} \rightarrow B^0)$	$< 1.1$	$\times 10^{-6}$	CL=90%
$\Gamma_7$	$\rho K^- K^- \times B(\bar{b} \rightarrow \Xi_b^-)$	$(3.7 \pm 0.8) \times 10^{-8}$		
$\Gamma_8$	$\rho K^- K^-$			
$\Gamma_9$	$\rho \pi^- \pi^-$			
$\Gamma_{10}$	$\rho K^- \pi^-$			
$\Gamma_{11}$	$\Lambda \pi^+ \pi^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$	$< 1.7$	$\times 10^{-6}$	CL=90%
$\Gamma_{12}$	$\Lambda K^- \pi^+ \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$	$< 8$	$\times 10^{-7}$	CL=90%
$\Gamma_{13}$	$\Lambda K^+ K^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$	$< 3$	$\times 10^{-7}$	CL=90%
$\Gamma_{14}$	$\Lambda_c^+ K^- \times B(\bar{b} \rightarrow \Xi_b^-)$	$(6 \pm 4) \times 10^{-7}$		
$\Gamma_{15}$	$\Lambda_b^0 \pi^- \times B(b \rightarrow \Xi_b^-)/B(b \rightarrow \Lambda_b^0)$	$(5.7 \pm 2.0) \times 10^{-4}$		
$\Gamma_{16}$	$\rho K^- \pi^+ \pi^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$	$(1.9 \pm 0.4) \times 10^{-6}$		
$\Gamma_{17}$	$\rho K^- K^- \pi^+ \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$	$(1.73 \pm 0.32) \times 10^{-6}$		
$\Gamma_{18}$	$\rho K^- K^+ K^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$	$(1.8 \pm 1.0) \times 10^{-7}$		

### $\Xi_b$ BRANCHING RATIOS

$\Gamma(\Xi^- \ell^- \bar{\nu}_\ell X \times B(\bar{b} \rightarrow \Xi_b^-))/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.9 \pm 1.2</math> OUR AVERAGE</b>	Error includes scale factor of 1.4.		
$3.0 \pm 1.0 \pm 0.3$	ABDALLAH	05C DLPH	$e^+ e^- \rightarrow Z^0$
$5.4 \pm 1.1 \pm 0.8$	BUSKULIC	96T ALEP	Excess $\Xi^- \ell^-$ over $\Xi^- \ell^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$5.9 \pm 2.1 \pm 1.0$	ABREU	95V DLPH	Repl. by ABDALLAH 05c

$\Gamma(J/\psi \Xi^- \times B(b \rightarrow \Xi_b^-))/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.102^{+0.026}_{-0.021}</math> OUR AVERAGE</b>			

$0.098^{+0.023}_{-0.016} \pm 0.014$  <sup>1</sup> AALTONEN 09AP CDF  $\rho \bar{p}$  at 1.96 TeV

$0.16 \pm 0.07 \pm 0.02$  <sup>2</sup> ABAZOV 07K D0  $\rho \bar{p}$  at 1.96 TeV

<sup>1</sup> AALTONEN 09AP reports  $[\Gamma(\Xi_b^- \rightarrow J/\psi \Xi^- \times B(b \rightarrow \Xi_b^-))/\Gamma_{\text{total}}] / [B(\Lambda_b^0 \rightarrow J/\psi(1S) \Lambda \times B(b \rightarrow \Lambda_b^0))] = 0.167^{+0.037}_{-0.025} \pm 0.012$  which we multiply by our best value  $B(\Lambda_b^0 \rightarrow J/\psi(1S) \Lambda \times B(b \rightarrow \Lambda_b^0)) = (5.8 \pm 0.8) \times 10^{-5}$ . Our first error is

their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABAZOV 07K reports  $[\Gamma(\Xi_b^- \rightarrow J/\psi \Xi^- \times B(b \rightarrow \Xi_b^-))/\Gamma_{\text{total}}] / [B(\Lambda_b^0 \rightarrow J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0))] = 0.28 \pm 0.09_{-0.08}^{+0.09}$  which we multiply by our best value  $B(\Lambda_b^0 \rightarrow J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)) = (5.8 \pm 0.8) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(J/\psi \Lambda K^- \times B(b \rightarrow \Xi_b^-))/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$**

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>2.45 \pm 0.19 \pm 0.35</math></b>	<sup>1,2</sup> AAIJ	17BE	LHCB $pp$ at 7 and 8 TeV

<sup>1</sup> AAIJ 17BE reports  $[\Gamma(\Xi_b^- \rightarrow J/\psi \Lambda K^- \times B(b \rightarrow \Xi_b^-))/\Gamma_{\text{total}}] / [B(\Lambda_b^0 \rightarrow J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0))] = (4.19 \pm 0.29 \pm 0.15) \times 10^{-2}$  which we multiply by our best value  $B(\Lambda_b^0 \rightarrow J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)) = (5.8 \pm 0.8) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Integrated over the  $b$ -baryon transverse momentum  $p_T < 25$  GeV and rapidity  $2.0 < y < 4.5$ .

**$\Gamma(\rho D^0 K^- \times B(\bar{b} \rightarrow \Xi_b^-))/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$**

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>(1.7 \pm 0.4 \pm 0.4) \times 10^{-6}</math></b>	<sup>1</sup> AAIJ	14H	LHCB $pp$ at 7 TeV

<sup>1</sup> AAIJ 14H reports  $[\Gamma(\Xi_b^- \rightarrow \rho D^0 K^- \times B(\bar{b} \rightarrow \Xi_b^-))/\Gamma_{\text{total}}] / [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\Lambda_b^0 \rightarrow \rho D^0 K^-)] = 0.44 \pm 0.09 \pm 0.06$  which we multiply by our best values  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ ,  $B(\Lambda_b^0 \rightarrow \rho D^0 K^-) = (4.6 \pm 0.8) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

**$\Gamma(\rho \bar{K}^0 \pi^- \times B(\bar{b} \rightarrow \Xi_b^-)/B(\bar{b} \rightarrow B^0))/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 1.6 \times 10^{-6}</math></b>	90	AAIJ	14Q	LHCB $pp$ at 7 TeV

**$\Gamma(\rho K^0 K^- \times B(\bar{b} \rightarrow \Xi_b^-)/B(\bar{b} \rightarrow B^0))/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 1.1 \times 10^{-6}</math></b>	90	AAIJ	14Q	LHCB $pp$ at 7 TeV

**$\Gamma(\rho K^- K^- \times B(\bar{b} \rightarrow \Xi_b^-))/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$**

VALUE (units $10^{-8}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>3.7 \pm 0.8 \pm 0.2</math></b>	<sup>1</sup> AAIJ	17F	LHCB $pp$ at 7, 8 TeV

<sup>1</sup> AAIJ 17F reports  $[\Gamma(\Xi_b^- \rightarrow \rho K^- K^- \times B(\bar{b} \rightarrow \Xi_b^-))/\Gamma_{\text{total}}] / [B(B^+ \rightarrow K^+ K^- K^+)] / [B(\bar{b} \rightarrow B^+)] = (2.65 \pm 0.35 \pm 0.47) \times 10^{-3}$  which we multiply by our best values  $B(B^+ \rightarrow K^+ K^- K^+) = (3.40 \pm 0.14) \times 10^{-5}$ ,  $B(\bar{b} \rightarrow B^+) = (40.8 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

**$\Gamma(\rho \pi^- \pi^-)/\Gamma(\rho K^- K^-)$   $\Gamma_9/\Gamma_8$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 0.56</math></b>	90	<sup>1</sup> AAIJ	17F	LHCB $pp$ at 7, 8 TeV

<sup>1</sup> Measures the ratio as  $0.28 \pm 0.16 \pm 0.13$ .

$\Gamma(\rho K^- \pi^-)/\Gamma(\rho K^- K^-)$   $\Gamma_{10}/\Gamma_8$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.98±0.27±0.09</b>	AAIJ	17F	LHCB <i>pp</i> at 7, 8 TeV

$\Gamma(\Lambda \pi^+ \pi^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}$   $\Gamma_{11}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.7 × 10<sup>-6</sup></b>	90	AAIJ	16W	LHCB <i>pp</i> at 7, 8 TeV

$\Gamma(\Lambda K^- \pi^+ \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}$   $\Gamma_{12}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.8 × 10<sup>-6</sup></b>	90	AAIJ	16W	LHCB <i>pp</i> at 7, 8 TeV

$\Gamma(\Lambda K^+ K^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}$   $\Gamma_{13}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.3 × 10<sup>-6</sup></b>	90	AAIJ	16W	LHCB <i>pp</i> at 7, 8 TeV

$\Gamma(\rho K^- \pi^+ \pi^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}$   $\Gamma_{16}/\Gamma$

VALUE (units 10 <sup>-6</sup> )	DOCUMENT ID	TECN	COMMENT
<b>1.91±0.35±0.18</b>	<sup>1</sup> AAIJ	18Q	LHCB <i>pp</i> at 7, 8 TeV

<sup>1</sup> AAIJ 18Q reports  $[\Gamma(\Xi_b \rightarrow \rho K^- \pi^+ \pi^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}] / [B(\Lambda_c^+ \rightarrow \rho K^- \pi^+)] / [B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] = (6.2 \pm 0.8 \pm 0.2 \pm 0.8) \times 10^{-3}$  which we multiply by our best values  $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (6.28 \pm 0.32) \times 10^{-2}$ ,  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = (4.9 \pm 0.4) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\rho K^- K^- \pi^+ \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}$   $\Gamma_{17}/\Gamma$

VALUE (units 10 <sup>-6</sup> )	DOCUMENT ID	TECN	COMMENT
<b>1.73±0.27±0.16</b>	<sup>1</sup> AAIJ	18Q	LHCB <i>pp</i> at 7, 8 TeV

<sup>1</sup> AAIJ 18Q reports  $[\Gamma(\Xi_b \rightarrow \rho K^- K^- \pi^+ \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}] / [B(\Lambda_c^+ \rightarrow \rho K^- \pi^+)] / [B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] = (5.6 \pm 0.6 \pm 0.4 \pm 0.5) \times 10^{-3}$  which we multiply by our best values  $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (6.28 \pm 0.32) \times 10^{-2}$ ,  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = (4.9 \pm 0.4) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\rho K^- K^+ K^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}$   $\Gamma_{18}/\Gamma$

VALUE (units 10 <sup>-6</sup> )	DOCUMENT ID	TECN	COMMENT
<b>0.18±0.09±0.02</b>	<sup>1,2</sup> AAIJ	18Q	LHCB <i>pp</i> at 7, 8 TeV

<sup>1</sup> AAIJ 18Q reports  $[\Gamma(\Xi_b \rightarrow \rho K^- K^+ K^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0))/\Gamma_{total}] / [B(\Lambda_c^+ \rightarrow \rho K^- \pi^+)] / [B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] = (0.57 \pm 0.28 \pm 0.08 \pm 0.10) \times 10^{-3}$  which we multiply by our best values  $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (6.28 \pm 0.32) \times 10^{-2}$ ,  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = (4.9 \pm 0.4) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>2</sup> AAIJ 18Q sees excess with a significance of 2.3 $\sigma$ . Using  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = (0.430 \pm 0.036) \times 10^{-2}$  and  $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (6.46 \pm 0.24) \times 10^{-2}$  the authors set two sided limit [0.11–0.25] at 90% C.L.

$$\Gamma(\Lambda_c^+ K^- \times B(\bar{b} \rightarrow \Xi_b)) / \Gamma(p D^0 K^- \times B(\bar{b} \rightarrow \Xi_b)) \quad \Gamma_{14}/\Gamma_4$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.36 ± 0.19 ± 0.02</b>	<sup>1</sup> AAIJ	14H LHCB	<i>pp</i> at 7 TeV

<sup>1</sup> AAIJ 14H reports  $[\Gamma(\Xi_b \rightarrow \Lambda_c^+ K^- \times B(\bar{b} \rightarrow \Xi_b)) / \Gamma(\Xi_b \rightarrow p D^0 K^- \times B(\bar{b} \rightarrow \Xi_b))] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+) / B(D^0 \rightarrow K^- \pi^+)] = 0.57 \pm 0.22 \pm 0.21$  which we multiply or divide by our best values  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.28 \pm 0.32) \times 10^{-2}$ ,  $B(D^0 \rightarrow K^- \pi^+) = (3.950 \pm 0.031) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

$$\Gamma(\Lambda_b^0 \pi^- \times B(b \rightarrow \Xi_b^-)) / B(b \rightarrow \Lambda_b^0) / \Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE (units 10 <sup>-4</sup> )	DOCUMENT ID	TECN	COMMENT
<b>5.7 ± 1.8<sup>+0.8</sup><sub>-0.9</sub></b>	<sup>1</sup> AAIJ	15BA LHCB	<i>pp</i> at 7, 8 TeV

<sup>1</sup> A signal is reported with a significance of 3.2 standard deviations in the decay chain of  $\Xi_b^- \rightarrow \Lambda_b^0 \pi^-$ ,  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ , and  $\Lambda_c^+ \rightarrow p K^- \pi^+$ .

## P AND CP VIOLATION

$$a_P(\Xi_b^0 \rightarrow p K^- K^- \pi^+)$$

Observable calculated as average of the triple products for  $\Xi_b^0$  and  $\bar{\Xi}_b^0$ , which is sensitive to parity violation.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-3.04 ± 5.19 ± 0.36</b>	<sup>1</sup> AAIJ	18AG LHCB	<i>pp</i> at 7, 8 TeV

<sup>1</sup> Measured over full phase space of the decay.

$$a_{CP}(\Xi_b^0 \rightarrow p K^- K^- \pi^+)$$

Observable calculated as half of the difference between triple products for  $\Xi_b^0$  and  $\bar{\Xi}_b^0$ , which is sensitive to *CP* violation.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-3.58 ± 5.19 ± 0.36</b>	<sup>1</sup> AAIJ	18AG LHCB	<i>pp</i> at 7, 8 TeV

<sup>1</sup> Measured over full phase space of the decay.

$$\Delta A_{CP}(\Xi_b^0 \rightarrow p K^- \pi^+ \pi^-)$$

$\Delta A_{CP} \equiv A_{CP}(\Xi_b^0 \rightarrow p K^- \pi^+ \pi^-) - A_{CP}(\bar{\Xi}_b^0 \rightarrow (\Xi_c^+ \rightarrow p K^- \pi^+) \pi^-)$

VALUE (units 10 <sup>-2</sup> )	DOCUMENT ID	TECN	COMMENT
<b>-17 ± 11 ± 1</b>	<sup>1</sup> AAIJ	19AH LHCB	<i>pp</i> at 7 and 8 TeV

<sup>1</sup> Full phase space.

$$\Delta A_{CP}(\Xi_b^0 \rightarrow p K^- \pi^+ K^-)$$

$\Delta A_{CP} \equiv A_{CP}(\Xi_b^0 \rightarrow p K^- \pi^+ K^-) - A_{CP}(\bar{\Xi}_b^0 \rightarrow (\Xi_c^+ \rightarrow p K^- \pi^+) \pi^-)$

VALUE (units 10 <sup>-2</sup> )	DOCUMENT ID	TECN	COMMENT
<b>-6.8 ± 8.0 ± 0.8</b>	<sup>1</sup> AAIJ	19AH LHCB	<i>pp</i> at 7 and 8 TeV

<sup>1</sup> Full phase space.



**$A_P(\Xi_b), \Xi_b^- - \Xi_b^+$  production asymmetry**

$$A_P(\Xi_b) = [\sigma(\Xi_b^-) - \sigma(\Xi_b^+)] / [\sigma(\Xi_b^-) + \sigma(\Xi_b^+)]$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>-2 ± 4 OUR AVERAGE</b>			
$1.1 \pm 5.6 \pm 1.9$	1,2 AAIJ	19AB LHCB	$pp$ at 7 and 8 TeV
$-3.9 \pm 4.9 \pm 2.5$	1,2 AAIJ	19AB LHCB	$pp$ at 13 TeV

<sup>1</sup> Baryon kinematic range  $p_T < 20$  GeV/c and  $2 < \eta < 6$ .

<sup>2</sup> Measured using previous measurements of  $A_P(\Lambda_b)$  in AAIJ 17BF.

 **$\Xi_b$  REFERENCES**

AAIJ	19AB PR D99 052006	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	19AH EPJ C79 745	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AG JHEP 1808 039	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18Q JHEP 1802 098	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BE PL B772 265	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BF PL B774 139	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BH PRL 119 181807	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BJ PRL 119 232001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17F PRL 118 071801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16W JHEP 1605 081	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15BA PRL 115 241801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14AA PRL 112 202001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BE NP B888 169	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BJ PRL 113 242002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14H PR D89 032001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14Q JHEP 1404 087	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14T PL B736 154	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14U PL B734 122	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14Z PRL 113 032001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14B PR D89 072014	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	13AV PRL 110 182001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	11X PRL 107 102001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09AP PR D80 072003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	07A PRL 99 052002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07K PRL 99 052001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	05C EPJ C44 299	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
BUSKULIC	96T PL B384 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	95V ZPHY C68 541	P. Abreu <i>et al.</i>	(DELPHI Collab.)